

AD-A049 152

NAVAL RESEARCH LAB WASHINGTON D C
HARMONIC GENERATION IN ND:LASER-PRODUCED PLASMAS.(U)
OCT 77 E A MCLEAN, J A STAMPER, B H RIPIN
NRL-MR-3626 SBIE-AD-E000 057

F/G 20/5

UNCLASSIFIED

1 OF 1
AD
A049152



END
DATE
FILMED
3-78
DDC

AD A049152

ad 000057

AD No. _____
DDC FILE COPY.

12

NRL Memorandum Report 3626

2

Harmonic Generation in Nd:Laser-Produced Plasmas

E. A. McLEAN, J. A. STAMPER, B. H. RIPIN,
J. M. McMAHON, and S. E. BODNER


*Laser Plasma Branch
Plasma Physics Division*

and

H. R. GRIEM

*University of Maryland
College Park, Maryland*

October 1977



NAVAL RESEARCH LABORATORY
Washington, D.C.

Approved for public release; distribution unlimited.

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER NRL Memorandum Report 3626	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) HARMONIC GENERATION IN Nd:LASER-PRODUCED PLASMAS.	5. TYPE OF REPORT & PERIOD COVERED Interim report, on a continuing NRL problem.	
7. AUTHOR(s) E. A. McLean, J. A. Stamper, B. H. Ripin, H. R. Griem*, J. M. McMahon and S. E. Bodner	6. PERFORMING ORG. REPORT NUMBER	
9. PERFORMING ORGANIZATION NAME AND ADDRESS Naval Research Laboratory Washington, D. C. 20375	8. CONTRACT OR GRANT NUMBER(s)	
11. CONTROLLING OFFICE NAME AND ADDRESS Energy Research and Development Administration Washington, D. C. 20545	10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NRL Problem H02-29A	
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) SBIE	12. REPORT DATE October 1977	
17. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. NRL-MR-3626	13. NUMBER OF PAGES 13	
18. SUPPLEMENTARY NOTES Sponsored by the Energy Research and Development Administration at the Naval Research Laboratory. *University of Maryland, College Park, Maryland	15. SECURITY CLASS. (of this report) UNCLASSIFIED	
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Laser harmonics Laser matter interaction Laser absorption Neodymium glass laser	16. DECLASSIFICATION/DOWNGRADING SCHEDULE	
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The fundamental and second through fifth harmonic spectral lines have been observed from the plasma produced when a 75-psec Nd:glass laser ($< 10^{16}$ W/cm ²) is focused onto a thick, planar polystyrene target. Both line profiles and relative intensities of these harmonics are given. approx. 10 to the 16th power W/cm ²		

DDC
RECEIVED
JAN 31 1978
B

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE
S/N 0102-014-6601

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

251 950

LB

HARMONIC GENERATION IN Nd:LASER-PRODUCED PLASMAS

There have been many observations of second harmonic light¹⁻⁸ emitted from laser-produced plasma. However, there have been no reported observations of third or higher harmonic emission in plasmas generated by a Nd:glass laser, although third harmonic light⁹⁻¹⁰ and higher harmonics¹¹ have been observed from CO₂ laser-produced plasmas. We report here the first observations of the third, fourth, and fifth harmonics lines produced when a high irradiance Nd:laser beam (1.06 μm) is focused onto a target. It is generally thought that integral harmonic emission is generated at or near the critical surface (where fundamental and local plasma frequencies are equal) and, therefore, can yield information on density gradients and electric fields in this region of the plasma.^{2,12} (A slower fall-off in the intensity of the harmonics would be an indication of a steeper density gradient or a higher field strength.) Thus, information on the harmonic emission is important in understanding the laser-target absorption process. These emissions may also interfere with diagnostic light probe studies such as shadowgraphy, interferometry, Faraday rotation, etc. which use light at the harmonic frequencies of the incident laser beam. As an example, a time-integrated photograph of third harmonic emitted light (3546 Å) taken through an interference filter (3546 \pm 50 Å) is shown in Fig. 1a. The small spatial extent

Note: Manuscript submitted September 28, 1977.

for		
White Section	<input checked="" type="checkbox"/>	
Buff Section	<input type="checkbox"/>	
D		
N		
D1		
DISTRIBUTION/AVAILABILITY CODES		
Dist.	AVAIL.	and/or SPECIAL
A		

of the emission ($\sim 5 \mu\text{m}$), even though averaged over time and spread by refraction effects, is indicative of a very localized emission region and, perhaps, of steep density gradients near the critical density.

The experimental arrangement is shown in Fig. 1b. One beam (6-10 J) from the NRL Pharos II Nd:laser ($1.06 \mu\text{m}$) operating at 75-psec pulse duration (FWHM) was focused at normal incidence onto a polystyrene (CH) slab target in an evacuated ($\leq 10^{-4}$ Torr) chamber using an f/1.9 aspheric lens. Prepulses were monitored on each shot and did not exceed $1 \mu\text{J}$ in these experiments. The half-energy focal diameter was measured¹³ to be $30 \mu\text{m}$ yielding an average irradiance of $\sim 10^{16} \text{ W/cm}^2$. A focal-shift monitor¹⁴ ensured that the target was in focus on each shot. The fractional absorption of laser light was inferred to be $\sim 50\%$ by measuring the light backscattered through the focusing lens and outside the lens with calorimeters.¹⁵ Simultaneous observations of the harmonic light were made with a 3/4-m Czerny-Turner mount spectrograph (resolution in first order $\sim 1 \text{ \AA}$ for a slit width of $100 \mu\text{m}$) and an interference filter - photovoltaic silicon photodiode combination (E.G.G. UV-100 B biased to $-22\frac{1}{2} \text{ V}$). These detectors viewed the target at 45° to the target normal and in a (horizontal) plane at 45° to the plane of the incident laser's electric field. The rise time of the photodiode and recording oscilloscope was $\sim 2 \text{ nsec}$, and the photodiode was proved linear in the range discussed here. Care was taken to block the higher order spectra from entering the spectrograph. An achromatic optical train was achieved by using

a spherical and a flat mirror to focus the plasma onto the spectrograph slit. Three types of film were used in the spectrograph to cover the range of wavelengths and sensitivities encountered in this experiment, i.e., Kodak I-Z plates, Kodak Royal-X Pan, and Polaroid Type 57. (All spectra required only one shot.)

Microdensitometer scans of representative time-integrated spectral profiles for the fundamental emission, ω_0 , and the integral harmonics through the fifth, $5\omega_0$, are shown in Fig. 2. The wavelength of each harmonic is indicated by a vertical dashed line. The peak of the fundamental line is blue-shifted by about 13 Å. If the shift (13 Å) is a Doppler shift, this would correspond to an average outward velocity of the reflecting surface of 3×10^7 cm/sec. A blue shift of the fundamental has also been observed by others.¹⁰ However, the centroids of the rest of the harmonic lines are red-shifted. It appears that the scattered fundamental line is produced by a different mechanism or spectral region than that which produces the harmonic lines. The $5\omega_0$ line at 2128 Å is just resolved above the plasma continuum. To observe this harmonic line, the spectrograph slit was opened to 500 μ m so that the resolution was reduced to ~ 5 Å. Because of the expected low signal level, we did not attempt to observe $6\omega_0$ at 1773 Å. All of the observed harmonic spectral widths are large compared to that of the incident laserwidth (~ 1 Å).

The spectral widths (FWHM) for the second, third, and fourth harmonics in wavelength and frequency units (to facilitate a comparison with theory) are shown in Table 1. It is seen that the spectral half-

widths in frequency units change at most by 60%, whereas the half-widths in wavelength units change by almost a factor of 3. The observed spectral broadening thus involves mechanisms other than Doppler broadening (either due to random or non-local mass motion) such as instabilities and magnetic fields.¹⁶

Assuming shot-to-shot reproducibility of the over 150 data shots, relative (total) intensities of the harmonics measured with the silicon photodiode-interference filter are shown in Fig. 3. These values of relative intensity are corrected for the transmission and the band pass of the filters and the spectral responsivity (given by the manufacturer) of the silicon photodiode. The error bars indicate the estimated error in the measurement due to the spectral responsivity uncertainty and the signal-to-noise ratio. There is a large decrease ($\sim 2 \times 10^2$) in the intensity of the second harmonic relative to the reflected or scattered fundamental; however, the next three harmonics decrease in steps of less than a factor of 10. These ratios of harmonic line intensities are the same (within a factor of 5 over a range of 5 orders of magnitude) as the corresponding harmonic intensities reported by the National Research Council Group¹¹ in Ottawa, Canada, using a CO₂ laser at an irradiance of $I = 10^{14} \text{ W/cm}^2$. This fact suggests that a comparable interaction takes place in Nd: and CO₂: laser irradiation which scales as $I\lambda^2$.

In summary, we report emission spectra and relative intensities of the fundamental and second through the fifth harmonics of the incident laser line when a Nd:laser (10^{16} W/cm^2) is focused onto a

CH slab target. The emission is sufficiently intense that the contrast in diagnostic probe studies at these harmonic wavelengths would be adversely affected. (This difficulty has been obviated at NRL by shifting the probe wavelength using a Raman cell to a nonharmonic wavelength¹⁷.) Although the present theories of optical absorption of intense light in plasma have not yet predicted the ratios, widths, and shifts at the higher harmonics, these harmonics should be directly related to the absorption process and would aid in estimating density gradients and electric fields produced in the absorbing layer. Hopefully, these experimental results will stimulate the appropriate theoretical explanations. An interesting result of this study is that the spectra and relative intensities of the harmonics produced in Nd:laser-target interactions at 10^{16} W/cm² are strikingly similar to recent results¹¹ found in CO₂ laser-produced plasmas at 10^{14} W/cm².

We would like to acknowledge the excellent technical assistance of E. Turbyfill, T. DeRieux, and A. Unzicker. Also, we would like to thank R. Decoste, R. H. Lehmberg, C. M. Armstrong, and F. C. Young for useful conversations and assistance and R. W. Whitlock for help in analyzing the data.

REFERENCES

- a. Work sponsored by the U.S. ERDA.
- b. Permanent address: Dept. of Physics and Astronomy, University of Maryland, College Park, MD.
1. B.H. Ripin, J.M. McMahon, E.A. McLean, W.M. Manheimer, and J.A. Stamper, Phys. Rev. Lett. 33, 634 (1974).
2. K. Eidmann and R. Sigel, Phys. Rev. Lett. 34, 799 (1975).
3. Ping Lee, D.V. Giovanielli, R.P. Godwin, and G.H. McCall, Appl. Phys. Lett. 24, 406 (1974).
4. J.L. Bobin, M. Decroisette, B. Meyer, and Y. Vitel, Phys. Rev. Lett. 30, 37 (1973).
5. C. Yamanaka, T. Yamanaka, T. Sasaki, J. Mizui, and H.B. Kang, Phys. Rev. Lett. 32, 1038 (1974).
6. A.A. Gorokhov, V.D. Dyatlov, R.N. Medvedev, A.D. Starikov, and V.G. Tuzov, ZhETF Pis. Red. 21, 111 (1975), [JETP Lett. 21, 49 (1975)]; and V.V. Alexander, S.I. Anisimov, M.V. Brenner, E.P. Velikov, V.D. Vikharev, V.P. Zotov, N.G. Kovalsky, M.I. Pergament and A.I. Yaroslavsky, ZhETF, 71, 1826 (1976).
7. T.A. Leonard, R.A. Cover, P. M. Campbell and R.A. Grandey, Conference Record - Abstracts IEEE 1977 International Conference on Plasma Science, May 23-25, 1977, Paper 4D12.
8. J. Soures, L.M. Goldman, and M. Lubin, Laser Interaction and Related Plasma Phenomena, H. Schwarz and H. Hora, Eds. (Plenum, New York 1974), Vol. 3B, p. 607.
9. H.A. Baldis, H. Pépin, and B. Grek, Appl. Phys. Lett. 27, 291 (1975).

10. E. Fabre, C. Garban, C. Popovics, A. Poquerusse, C. Stenz, and J. Virmont, Proceedings of IAEA 6th International Conference on Plasma Physics and Controlled Nuclear Fusion Research, Berchtesgaden, West Germany, Oct. 6-13, 1976, Paper No. IAEA-CN 35/G 3-4.
11. N. H. Burnett, H. A. Baldis, M. C. Richardson, and G. D. Enright, Appl. Phys. Lett. 31, 172 (1977).
12. R. E. Kidder, Laser Interaction with Matter, edited by C. Yamanaka (Japan Society for Promotion of Science, Tokyo, 1973), p. 331.
13. B. H. Ripin, NRL Memo Report 3315, 1976 (unpublished).
14. J. A. Stamper, Appl. Optics 51, 2020 (1976).
15. B. H. Ripin, F. C. Young, R. Decoste, C. M. Armstrong, E. A. McLean, and J. A. Stamper, to be published.
16. W. L. Kruer and K. G. Estabrook, Laser Interactions and Related Plasma Phenomena, H. Schwarz and H. Hora, Eds. (Plenum, New York, 1976).
17. R. H. Lehmberg, to be published.

Table 1 — Spectral half-widths (FWHM) of the harmonics in frequency and wavelength units. The estimated error from the photographic plate calibration and a small number of representative shots is $\pm 10\%$.

HARMONIC	$\Delta\omega_{FWHM}$ [rad sec ⁻¹]	$\Delta\lambda_{FWHM}$ [Å]
$2\omega_0$ (5320 Å)	1.9×10^{13}	29
$3\omega_0$ (3547 Å)	1.7×10^{13}	11
$4\omega_0$ (2660 Å)	2.7×10^{13}	10





Fig. 1 — a) A time-integrated photograph of the third harmonic light (3546 \AA) emitted tangential to the target surface when a 75-psec, $\sim 10^{16} \text{ W/cm}^2$ laser pulse strikes a CH_2 plane target. The laser beam was incident from the left. A 3547 \AA interference filter with a bandwidth of 50 \AA was used in front of the camera.
b) The experimental arrangement.




Fig. 2 — Representative time-integrated spectral line profiles are shown for the fundamental through the fifth harmonic. This radiation was observed at 45 degrees from the incident laser beam. Extraneous marks are indicated on the ω_0 and $4\omega_0$ profiles. The calculated wavelength of the center of each spectral line is indicated with a dashed line.




Fig. 3 — Intensities of the harmonics relative to the fundamental emission at 45° to the incident laser beam. The error bars are estimated from the spectral responsivity uncertainty and the signal-to-noise ratio.